

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US05/004933

International filing date: 17 February 2005 (17.02.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/546,559
Filing date: 20 February 2004 (20.02.2004)

Date of receipt at the International Bureau: 17 March 2005 (17.03.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1293665

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

March 08, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/546,559

FILING DATE: February 20, 2004

RELATED PCT APPLICATION NUMBER: PCT/US05/04933



Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

13049 U.S. PTO
022004

PTO/SB/16 (01-04)
Approved for use through 07/31/2006. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. ER 426254048 US

U.S. PTO
60/546559
022004

INVENTOR(S)					
Given Name (first and middle [if any])		Family Name or Surname		Residence (City and either State or Foreign Country)	
Ladislav Vlastislav		Cvak Mojczek		Czech Republic Czech Republic	
Additional inventors are being named on the _____ separately numbered sheets attached hereto					
TITLE OF THE INVENTION (500 characters max)					
PROCESS FOR THE MANUFACTURE OF LYSERGIC ACID					
Direct all correspondence to: CORRESPONDENCE ADDRESS					
<input type="checkbox"/> Customer Number: _____					
OR					
<input checked="" type="checkbox"/> Firm or Individual Name		Dennis A. Emma, Ph.D.			
Address		IVAX CORPORATION			
Address		4400 Biscayne Boulevard			
City		State	Florida	Zip	33137
Country		Telephone	305 575 6075	Fax	305 575 6064
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification Number of Pages 11		<input type="checkbox"/> CD(s), Number _____			
<input type="checkbox"/> Drawing(s) Number of Sheets _____		<input type="checkbox"/> Other (specify) _____			
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76					
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT					
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.				FILING FEE Amount (\$)	
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees.				\$160.00	
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: 50-0943					
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.					
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.					
<input checked="" type="checkbox"/> No.					
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____					

Respectfully submitted,

[Page 1 of 2]

Date February 20, 2004

SIGNATURE Dennis A. Emma

REGISTRATION NO. 50980

TYPED or PRINTED NAME Dennis A. Emma, Ph.D.

(if appropriate)

Docket Number: GAL0020-P-USA

TELEPHONE 305 575 6075

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE


In re Application of: Ladislav Cvak and Vlastislav Mojczek
Serial Number: To be assigned
Filing Date: HEREWITH
Title: PROCESS FOR THE MANUFACTURE OF LYSERGIC ACID
Attorney Docket No.: GAL0019-P-USA

CERTIFICATION UNDER 37 C.F.R. § 1.10

I hear by certify that the attached papers are being deposited with the United States Postal Service as "Express Mail Post Office to Addressee" Service of the United States Postal Service (UPS) under 37 C.F.R. 1.10 on February 20, 2004, and is addressed to: MAIL STOP PROVISIONAL PATENT APPLICATION, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450

ER 426254048 US

"Express Mail Label No."


Angelo J. Mignanelli

MAIL STOP PROVISIONAL PATENT APPLICATION

Commissioner for Patents
P.O. BOX 1450
Alexandria, VA 22313-1450

Dear Sir:

Enclosed herewith for filing with the United States Patent and Trademark Office in the above-identified Provisional Patent Application pursuant to 37 C.F.R. § 1.53 (c) are the following documents:

1. Provisional Application Cover Sheet (one page);
2. Provisional Patent Application (11 pages); and
3. Return Postcard.

Respectfully Submitted,



Dennis A. Emma, Ph.D.
Registration No. 50,980
Attorney for Applicants

IVAX CORPORATION
4400 Biscayne Boulevard
Miami, FL 33137
Tel.: (305) 575-6075
Fax: (305) 575-6064

Date: February 20, 2004

PROCESS FOR THE MANUFACTURE OF LYSERGIC ACID

Inventors: Ladislav Cvak and Vlastislav Mojczek
(Attorney Docket Number GAL0019-P-USA)

FIELD OF THE INVENTION




[0001] The present invention relates to a process for the manufacture of lysergic acid by isomerizing paspalic acid in high yield.

BACKGROUND OF THE INVENTION

[0002] Lysergic acid is a basic structural part of natural ergot alkaloids. It is manufactured in large scale as an intermediate for the synthesis of several semisynthetic ergot alkaloids, which have found uses as drugs, *e.g.*, ergometrine, methylergometrine, methysergide and nicergoline.

[0003] Paspalic acid is also an ergot alkaloid, which is readily available by fermentation. The conversion of paspalic acid to lysergic acid is known by various methods. The process, however, results in the formation of unwanted impurities such as the epimer, isolysergic acid, when known isomerization techniques are used.

[0004] Lysergic acid, paspalic acid and isolysergic acid are natural chiral compounds with the *R* configuration on the chiral center in position 5 of their skeleton. As used herein, lysergic acid, isolysergic acid and paspalic acid mean d-lysergic acid, d-isolysergic acid and d-paspalic acid. Their structures are depicted below.



[0005] Lysergic acid is typically manufactured by hydrolyzing natural ergot alkaloids like ergotamine or ergotoxine isolated from ergot. Another typical preparation involves the partial synthesis from natural precursors which are available by fermentation, such as by the hydrolysis of lysergic acid hydroxy-ethylamide or the isomerization of paspalic acid.

[0006] Paspalic acid and its isomerization to lysergic acid was first described by Kobel (Helv. Chim. Acta 47:1052 (1964)). The isomerization was accomplished by boiling paspalic acid in diluted aqueous sodium hydroxide. When the isomerization of paspalic acid was carried out by this procedure, however, the conversion to lysergic acid was unsatisfactory (more than 5% of paspalic acid remained in the reaction mixture) and the high temperature used caused significant decomposition of the product. Similar results were obtained with other procedures described by Kobel *at al.* (Helv. Chim. Acta 64:478 (1981) and JP 70013302), both using isomerization in boiling aqueous potassium hydroxide. Moreover, the reaction mixture contained a significant amount of isolysergic acid (more than 25 %) which decreased both the yield and the quality of the resulting lysergic acid (the isolated lysergic acid contained about 5 % of isolysergic acid). Therefore further purification of the product was necessary. Another drawback was the low concentration of paspalic acid in the reaction mixture, which in turn required a large volume reactor.

[0007] Recently a new process of preparing lysergic acid was disclosed which described the use of aqueous tetraalkylammonium hydroxides to isomerize paspalic acid. (U.S. Patent No. 6,242,603). The reported yield of isolated lysergic acid containing about 3 % of isolysergic acid was about 80 %. The content of paspalic acid in the product was not disclosed. This reference also described the preparation of lysergic acid by above-mentioned process including isomerization in boiling diluted sodium or potassium hydroxide. These comparative examples describe the formation of lysergic acid in less than 60% yields.

SUMMARY OF THE DISCLOSURE

[0008] An advantage of the present invention is a facile method of preparing lysergic acid from paspalic acid in high yields and quality.

[0009] These and other advantages are satisfied, at least in part, by a process for the manufacture of lysergic acid which includes isomerization of paspalic acid by a concentrated

aqueous solution of sodium or potassium hydroxide. The process can advantageously be carried out within a few hours and at relatively low temperature. Isolation of essentially pure lysergic acid can be achieved in yields of greater than 70 % and containing less than about 1 wt % of paspalic acid and less than about 1 wt % of isolysergic acid. The process can additionally provide epimerization of any unwanted isolysergic acid to the desired lysergic acid.

[0010] Embodiments of the present invention include combining paspalic acid, water and a metal hydroxide, *e.g.*, potassium or sodium hydroxide, for a sufficient time to form lysergic acid, wherein the amount of the paspalic acid and metal hydroxide is in sufficient quantity to cause a phase separated reaction mixture. Additional embodiments include acidifying the reaction mixture after isomerization to form a lysergic acid salt; extracting lysergic acid from the salt with a mixture of methanol and aqueous ammonia; partial evaporation of the extract, crystallizing the extracted lysergic acid in a crystallization medium; separating the crystalline lysergic acid from its crystallization medium; washing the crystalline lysergic acid with water and methanol; and combining the crystallization medium and the washing solvent with additional paspalic acid, water, and a metal hydroxide to manufacture additional lysergic acid in high yield and quality.

[0011] Additional advantages of the present invention will become readily apparent to those skilled in this art from the following detailed description, wherein only the preferred embodiment of the invention is shown and described, simply by way of illustration of the best mode contemplated of carrying out the invention. As will be realized, the invention is capable of other and different embodiments, and its several details are capable of modifications in various obvious respects, all without departing from the invention. Accordingly, the descriptions are to be regarded as illustrative in nature, and not as restrictive.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0012] After experimentation and investigation, it was discovered that paspalic acid can be isomerized to lysergic acid by concentrated aqueous solutions of metal hydroxides in surprisingly high yield and quality. In particular, it was discovered that the use of

concentrated aqueous solutions of a metal hydroxide, e.g., sodium or potassium hydroxide, had positive effects on both the yield and the quality of the formed lysergic acid. Both, the yield and quality of the isolated lysergic acid were substantially better than that known to the art e.g. described in U. S. Patent 6,242,603 B1.

[0013] Based upon experimentation, it was discovered that when a sufficient amount of paspalic acid is dissolved in least about a 12 weight percent aqueous sodium or potassium hydroxide solution, a phase separated reaction mixture forms. Without being bound by any theory, it is believed that the phase separated reaction mixture provides an excellent medium for isomerizing of paspalic acid. Paspalic acid can be converted to lysergic acid in such phase separated medium under relatively mild conditions, e.g., after about 4 hours of mixing at a temperature of about 50 °C. Under these reaction conditions, the conversion of paspalic acid was greater than 98 % so that the reaction mixture contained less than about 2.0 % of paspalic acid and a balance concentration of isolysergic acid was about 18 % (both determined by HPLC).

[0014] In general, the present invention is practiced by combining paspalic acid, water and a metal hydroxide. Because of convenience, the paspalic acid is typically added to the aqueous solution of the metal hydroxide. In one aspect of the present invention, the ingredients, paspalic acid, water, and a metal hydroxide are combined in such quantity that a phase separated reaction mixture is formed. The ingredients are combined and, optionally, agitated for a time sufficient to achieve high conversion to lysergic acid. The temperature of the mixture can also be increased to the reaction rate up to the reflux temperature nevertheless the temperature between about 40 °C to about 60 °C was found out to be optimal. The temperature above this range can also deleteriously decompose the formed lysergic acid.

[0015] After formation, lysergic acid can be separated from the reaction mixture. In one embodiment for practicing the present invention, lysergic acid is separated from the reaction mixture by acidification, e.g., with sulfuric acid, to form a salt, which is more readily separated from the reaction mixture. For example, acidifying the reaction mixture to a pH below about 4.0 with sulphuric acid forms lysergic acid sulfate salt which precipitates from the reaction mixture and can then be readily collected by filtration. Experiments have shown that the crude salt contains roughly the same proportion of lysergic acid and isolysergic acid

formed by isomerizing of paspalic acid. Hence, it may be desired to further isolate the formed lysergic acid from its epimer.

[0016] In another aspect of the present invention, the process includes dissolving the crude lysergic acid salt in a mixture of methanol and aqueous ammonia to regenerate lysergic acid from its salt. This can be followed by crystallization of the lysergic acid which can be advantageously carried out by evaporating the ammonia and partial evaporation of the methanol from the solution and cooling the solution. After separating the crystalline lysergic acid from its crystallization medium in this example, it was found that the obtained crystalline lysergic acid contained less than about 10% of isolysergic acid. The content of this epimer can be further decreased by washing the crystalline lysergic acid with methanol.

[0017] In practising another aspect of the present invention, lysergic acid containing isolysergic acid impurity can be purified by washing crystalline lysergic acid with methanol to reduce the amount of isolysergic acid impurity to less than about 3 wt%. In practice, lysergic acid containing less than 1% of isolysergic acid can be finally obtained. It is believed that isolysergic acid is more readily dissolved in methanol than lysergic acid and that washing the desired product with methanol provides a convenient way to further purify lysergic acid.

[0018] Experimentation has shown that the mother liquors from the crystallization and washing steps contain mainly isolysergic acid. These solutions can be returned back to the process of isomerization, where the isolysergic acid is epimerized to the desired lysergic acid thus increasing the overall yield of a multi-batch or continuous process. This process, including the recycling of the mother liquors, was repeated up to ten times without any problems with the quality of the obtained product. Total yield of a multi-batch process, including the recycling of the mother liquors containing isolysergic acid, was up to about 90% and the quality of obtained lysergic acid was very high. For example, the average content of the main impurities, paspalic acid and isolysergic acid, was below about 1% individually.

[0019] Through experimentation it was found, no significant difference between the use of sodium or potassium hydroxides. Both hydroxides form a two-phase system when at least about one weight part of paspalic acid is dissolved in no more than about 20 weight parts of an aqueous solution containing at least about one weight part of sodium or potassium

hydroxides in 8 weight parts of water. Other metal hydroxides are believed useable in similar amounts, nevertheless because the use of other hydroxides is lacking the practical sense, it was not further investigated.

[0020] The following examples illustrate, but do not limit, the process. While the first example describes the first isomerization of paspalic acid so that it did not include any recycled material, the second example describes a typical manufacturing process where the mother liquors from a previous batch are returned to the next batch.

EXAMPLES

Example 1

The preparation of lysergic acid by a process which does not include recycling the mother liquors after crystallization of lysergic acid was achieved as follows:

100.0 g of paspalic acid (titration assay 98.5 %) was dissolved in 1000 ml of 5 % aqueous sodium hydroxide and then 150 g of sodium hydroxide was added to the solution. A formation of two-phase system was observed. The obtained two-phase system was then mixed for about 4 hours at about 50 °C under nitrogen. Then the reaction mixture was diluted with 1000 ml of water, cooled to 10 °C and acidified to a pH of about 3.5 with 40% sulfuric acid. A suspension of crystalline lysergic acid sulfate formed was then mixed for about 2 hours at about 5 °C. Then the crystalline lysergic acid sulfate was filtered off. The lysergic acid sulfate was extracted three times with a 500 ml of mixture of methanol and aqueous ammonia 95:5 (v/v) and the joined extracts were evaporated to about 200 grams, diluted with 200 ml water and left to crystallize at about 5 °C for 24 hours. The crystalline lysergic acid was then separated and washed by about 100 ml of water and three times with about 100 ml of methanol. After vacuum drying (24 hours at 60 °C and 30 mbar), 73.4 grams of lysergic acid was obtained (titration assay 99.1 %, content of paspalic acid 0.5 %, content of isolysergic acid 0.8 %). The mother liquors after crystallization of lysergic acid and the methanol solution obtained after washing the crystalline product were evaporated to a volume of about 200 ml and were then added to the next batch, Example 2.

Example 2

The manufacture of lysergic acid in greater than about 90% yield was achieved by adding the mother liquors from the previous batch (Example 1) to the isomerisation process as follows: About 50 grams of sodium hydroxide was dissolved in about 800 ml of water and 200 ml of the concentrated mother liquors from the Example 1. 100.0 grams of paspalic acid (titration assay 98.5 %) and finally 150 g of sodium hydroxide were added to the solution. A two-phase reaction mixture was subsequently formed. The obtained two-phase system was mixed for about 4 hours at about 50 °C under nitrogen. Then the reaction mixture was diluted with 1000 ml of water, cooled to 10 °C and acidified to about pH 3.5 with 40% sulfuric acid. The obtained suspension was mixed for 2 hours at about 5 °C and then crystalline lysergic acid sulfate was filtered off. The lysergic acid sulfate was extracted three times with 500 ml of mixture of methanol and aqueous ammonia 95:5 (v/v) and the joined extracts were evaporated to about 200 grams, diluted with 200 ml water and let to crystallize at about 5 °C for 24 hours. The crystalline lysergic acid was then separated and washed by 100 ml of water and three times with 100 ml of methanol. After vacuum drying (24 hours at 60 °C and 30 mbar), 90.8 grams of lysergic acid was obtained (titration assay 98.7 %, content of paspalic acid 0.6 %, content of isolysergic acid 0.9 %). The mother liquors after crystallization of lysergic acid and the methanol solution obtained after washing of the crystalline product were evaporated to a volume of about 200 ml and were added to the next batch.

In this disclosure there is described only the preferred embodiments of the invention and but a few examples of its versatility. It is to be understood that the invention is capable of use in various other combinations and environments and is capable of changes or modifications within the scope of the inventive concept as expressed herein.

What is Claimed Is:

1. A process for the manufacture of lysergic acid by isomerization of paspalic acid, in which the isomerization is brought about by a phase separated system comprising paspalic acid and an aqueous solution of sodium hydroxide or potassium hydroxide.
2. A process according to claim 1, in which at least about one weight part of paspalic acid is dissolved in about 20 weight parts of the aqueous solution of sodium hydroxide or potassium hydroxide.
3. A process according to claims 1 and 2, in which the aqueous solution of sodium hydroxide or potassium hydroxide is prepared by dissolving at least about one weight part of sodium hydroxide or potassium hydroxide in about eight parts of water.
4. A process according to claim 1, in which the isomerization is performed at the temperature in the range from about 20 °C to about 100 °C.
5. A process according to claim 1, in which the isomerization is performed at the temperature from about 40 °C to about 60°C.
6. The process of claim 1 wherein the lysergic acid is manufactured in a yield of no less than 70%.
7. A process according to claim 1, comprising crystallizing the lysergic acid from water or methanol or their mixtures and recycling the mother liquors back to the isomerization process.

8. A process for the manufacture of lysergic acid, the process comprising combining paspalic acid, water, and a metal hydroxide to manufacture lysergic acid, wherein the paspalic acid and metal hydroxide are in sufficient quantity to cause a phase separated reaction mixture.
9. The process of claim 8 further comprising acidifying the reaction mixture after isomerization to form crystalline salt of lysergic acid.
10. The process of the claim 9 wherein the acidification is accomplished by sulphuric acid.
11. The process of the claim 9 further comprising extracting lysergic acid from the lysergic acid salt by a mixture of methanol and aqueous ammonia to form a lysergic acid solution.
12. The process of the claim 11 further comprising partial evaporating of the lysergic acid solution and crystallizing lysergic acid.
13. The process of the claim 12 further comprising separating the crystalline lysergic acid from the crystallization medium and washing the crystalline product with methanol.
14. The process of claim 13 further comprising combining the crystallization medium and washing methanol with additional paspalic acid, water, and a metal hydroxide to form additional lysergic acid.
15. The process of claim 13 wherein the separated crystallized lysergic acid contains less than about 1 wt % of paspalic acid and less than about 1 wt % of isolysergic acid.

16. A process of purifying lysergic acid, containing isolysergic acid impurity, the process comprising washing crystalline lysergic acid with methanol to reduce the amount of isolysergic acid impurity to less than about 3 wt %.
17. The process of claim 16 comprising washing crystalline lysergic acid with methanol to reduce the amount of lysergic acid impurity to less than about 1 wt %.

ABSTRACT

Lysergic acid is formed in high yields and quality by isomerizing paspalic acid in a phase separated system formed by paspalic acid and concentrated aqueous metal hydroxide solution.